

	Family	Species	Host	
wasp	Cynipidae	<i>Andricus atrimentus</i>	<i>Q. douglasii</i>	
wasp	Cynipidae	<i>Andricus chrysolepidicola</i>	<i>Q. douglasii</i>	
wasp	Cynipidae	<i>Andricus fullawayi</i>	<i>Q. lobata</i>	
wasp	Cynipidae	<i>Andricus gigas</i>	<i>Q. douglasii</i>	
wasp	Cynipidae	<i>Andricus kingi</i>	<i>Q. lobata</i>	
wasp	Cynipidae	<i>Andricus lasius</i>	<i>Q. chrysolepis</i>	
wasp	Cynipidae	<i>Andricus parmula</i>	<i>Q. douglasii</i>	
wasp	Cynipidae	<i>Andricus quercuscalifornicus</i>	<i>Q. lobata</i>	
wasp	Cynipidae	<i>Andricus reniformis</i>	<i>Q. chrysolepis</i>	
wasp	Cynipidae	<i>Andricus spectabilis</i>	<i>Q. chrysolepis</i>	
wasp	Cynipidae	<i>Andricus wiltzae</i>	<i>Q. lobata</i>	
wasp	Cynipidae	<i>Antron quercusechinus</i>	<i>Q. douglasii</i>	
wasp	Cynipidae	<i>Besbicus conspicuus</i>	<i>Q. douglasii</i>	
wasp	Cynipidae	<i>Besbicus multipunctatus</i>	<i>Q. douglasii</i>	
wasp	Cynipidae	<i>Callyritis perdens</i>	<i>Q. kelloggii</i>	
wasp	Cynipidae	<i>Dischalcaspis canescens</i>	<i>Q. douglasii</i>	
wasp	Cynipidae	<i>Dischalcaspis chrysolepidis</i>	<i>Q. chrysolepis</i>	
wasp	Cynipidae	<i>Dryocosmus asymmetricus?</i>	<i>Q. chrysolepis</i>	
wasp	Cynipidae	<i>Heteroecus flavens</i>	<i>Q. chrysolepis</i>	
wasp	Cynipidae	<i>Heteroecus melanoderma</i>	<i>Q. chrysolepis</i>	growth form altered by inquilines or parasitoids
wasp	Cynipidae	<i>Heteroecus pacificus</i>	<i>Q. chrysolepis</i>	unisexual and bisexual galls
wasp	Cynipidae	<i>Heteroecus sanctaclarae</i>	<i>Q. chrysolepis</i>	
wasp	Cynipidae	<i>Liodora pattersonnae</i>	<i>Q. douglasii</i>	
wasp	Cynipidae	<i>Neuroterus saltatorius</i>	<i>Q. douglasii</i>	
wasp	Cynipidae	<i>Paracraspis guadaloupensis</i>	<i>Q. chrysolepis</i>	
wasp	Cynipidae	spring leaf gall (undescribed species or alternate generation of described species, in Russo)	<i>Q. lobata</i>	
wasp	Cynipidae	Club Vein Gall (undescribed species pictured in Russo)	<i>Q. chrysolepis</i>	
midge	Cecidomyiidae	either " <i>Dasineura</i> " silvestrii (Russo p185) or an undescribed species (Russo p186)	<i>Q. chrysolepis</i>	folded leaf gall(s)
aphid	Aphididae	<i>Pemphigus populivenae</i>	<i>Populus fremontii</i>	cottonwood folded leaf galls -- aphids inside
aphid	Aphididae	<i>Pemphigus populitransversus</i>	<i>Populus fremontii</i>	cottonwood petiole gall
midge	Cecidomyiidae	<i>Rhopalomyia californica</i>	<i>Baccharis pilularis</i>	coyote bush midge gall
midge	Cecidomyiidae	<i>Rhopalomyia baccharis</i>	<i>Baccharis pilularis</i>	coyote bush stem gall (midge)
moth	Gelechiidae	<i>Gnorimoschema baccharisella</i>	<i>Baccharis pilularis</i>	coyote bush moth stem gall
mite	Eriophyidae	<i>Aculops tetanothrix</i>	<i>Salix sp.</i>	red mite galls
midge	Cecidomyiidae	<i>Rabdophaga salicisbrassicoides</i>	<i>Salix sp.</i>	rosette gall
		??	<i>Salix sp.</i>	folded leaf gall
midge	Cecidomyiidae	<i>Iteomyia sp. A</i>	<i>Salix sp.</i>	leaf gall (teeth)
sawfly (wasp)	Tenthredinidae	<i>Pontania sp.</i>	<i>Salix sp.</i>	Pontania leaf gall
midge or sawfly	Cecidomyiidae or Tenthredinidae	<i>Rabdophaga sp. or Euura sp.</i>	<i>Salix sp.</i>	stem swelling gall -- need to cut open for ID

listed compiled by Joyce Gross, 12/2011
joyceg@berkeley.edu
Blue Oak Ranch Reserve